

Shrimp Maturation

Maturation Conditioning Diet



Shrimp Maturation Conditioner is formulated as an integral part of the dietary needs of broodstock. The diet utilizes high quality ingredients such as marine proteins and oils to help broodstock satisfy their nutritional requirements. Shrimp Maturation Conditioner helps eliminate the normal lag time between ablation and initial ovary maturation in female penaeid shrimp. When the diet is fed to broodstock from approximately 20 grams to harvest, the normal lag time associated with ovarian maturation is virtually eliminated.

- Quicker growth of broodstock.
- Promotes rapid ovary development in maturation systems.
- Larger spawns and increased mating in maturation tanks.
- Rapid growth allows increased generations in genetic breeding program.
- Can be used with automatic feeders to optimize growth and production of broodstock.
- Contains **Vpak** (Vitality Pak) to enhance disease resistance.



Shrimp Maturation

Disclaimer: Photos of feed may vary from screen appearance, printing and lot production. This is a visual to give you a representation of particle range and color approximate. Consult your representative for a sample today.

GUARANTEED ANALYSIS

Crude Protein	Min 40.0%	Moisture	Max 10.0%
Crude Fat	Min 9.0%	Ash	Max 13.0%
Crude Fiber	Max 3.0%	Phosphorus	Min 1.1%

PARTICLE SIZE & PACKAGING

3/32" (2.4mm) pellet

Nitrogen flushed bags: 9.07 kg (20 lb) 2 per case.

INGREDIENTS

A nutritionally balanced highly digestible formulation of animal protein products of marine origin, plant protein (including algae), yeast, fish and vegetable oils, vegetable starches, vitamin and mineral premixes, anti-oxidants, pigments, and biodegradable binders.

STORAGE INSTRUCTIONS

- Poly-woven bag: best if used within twenty-four (24) months from date of manufacture.
- Store in a cool (22° C/72° F), dry, well-ventilated area away from sunlight.
- Refrigeration or frozen storage will extend shelf life.
- Rotate stock to use the old feed first ("first in, first out" principle.)

PRODUCT APPLICATION

- Feed directly to broodstock rearing ponds or raceways similar to normal growout feed.
- Feed ration of 3% biomass per day (dry weight basis) starting at 20 grams to harvest.
- Consult your area Zeigler technical specialist for additional feed guidance.

Copyright © 2022 by Zeigler Bros., Inc.

All rights reserved. No part of this publication may be reproduced, distributed, or transmitted in any form or by any means, including photocopying, recording, or other electronic or mechanical methods, without the prior written permission of the publisher, except in the case of brief quotations embodied in critical reviews and certain other non-commercial uses permitted by copyright law. For permission requests, write to the publisher, addressed "info@zeiglerfeed.com."



Zeigler Bros., Inc.
400 Gardners Station Road
Gardners, PA 17324 USA

www.zeiglerfeed.com
info@zeiglerfeed.com

717-677-6181
717-677-6826 fax

GLOBAL G.A.P.
The Global Standard for Safe and Sustainable Agriculture

